



# CHECKLIST FOR PRESCRIBING OPIOIDS FOR CHRONIC PAIN

For primary care providers treating adults (18+) with chronic pain ≥ 3 months, excluding cancer, palliative, and end-of-life care

## When CONSIDERING long-term opioid therapy

- Set realistic goals for pain and function based on diagnosis (eg, walk around the block).
- Check that non-opioid therapies tried and optimized.
- Discuss benefits and risks (eg, addiction, overdose) with patient.
- Evaluate risk of harm or misuse.
  - Discuss risk factors with patient.
  - Check prescription drug monitoring (CURES) data.
  - Check urine drug screen.
- Set criteria for stopping or continuing opioids.
- Assess baseline pain and function (daily activities).
- Schedule initial reassessment within 1-4 weeks.
- Prescribe short-acting opioids using lowest dosage on product labeling; match duration to scheduled reassessment.

## When REASSESSING at return visit

Continue opioids only after confirming clinically meaningful improvements in pain and function without significant risks or harm.

- Assess pain and function; compare results to baseline.
- Evaluate risk of harm or misuse:
  - Observe patient for signs of over-sedation or overdose risk.
    - If yes: Taper dose.
  - Check CURES.
  - Check for opioid use disorder if indicated (eg, difficulty controlling use).
    - If yes: Refer for treatment.
- Check that non-opioid therapies optimized.
- Determine whether to continue, adjust, taper, or stop opioids.
- Calculate opioid dosage morphine milligram equivalent (MED)
  - If ≥50 MED/day total (≥ 50 mg hydrocodone; ≥ 30 mg oxycodone) increase frequency of follow-up; consider offering naloxone.
  - Avoid ≥90 MED/day total (≥90 mg hydrocodone; ≥60 mg oxycodone), or carefully justify; consider specialist referral.
- Schedule reassessment at monthly intervals.

## Clinical Reminders

- Opioids are not first-line or routine therapy for chronic pain.
- Establish and measure goals for pain and function.
- Discuss benefits and risks and availability of nonopioid therapies with patient.
- Use immediate-release opioids when starting.
- Start low and go slow.
- When opioids are needed for acute pain, prescribe no more than needed.
- Do not prescribe ER/LA opioids for acute pain.
- Follow-up and re-evaluate risk of harm; reduce dose or taper and discontinue if needed.
- Evaluate risk factors for opioid related harms.
- Check CURES for high dosages and prescriptions from other providers.
- Use urine drug testing to identify prescribed substances and undisclosed use.
- Avoid concurrent benzodiazepine and opioid prescribing.
- Arrange treatment for opioid use disorder if needed.

### EVIDENCE ABOUT OPIOID THERAPY

- Benefits of long-term opioid therapy for chronic pain not well supported by evidence.
- Short-term benefits small to moderate for pain; inconsistent for function.
- Insufficient evidence for long-term benefits in low back pain, headache, and fibromyalgia.

### NON-OPIOID THERAPIES

Use alone or combined with opioids, as indicated:

- Non-opioid medications (eg, NSAIDs, TCAs, SNRIs, anti-convulsants).
- Physical treatments (eg exercise therapy, weight loss).
- Behavioral treatment (eg, CBT).
- Procedures (eg, intra-articular corticosteroids).

### EVALUATING RISK OF HARM OR MISUSE

Known risk factors include:

- Illegal drug use; prescription drug use for nonmedical reasons.
- History of substance use disorder or overdose.
- Mental health conditions (eg, depression, anxiety).
- Sleep-disordered breathing.
- Concurrent benzodiazepine use.

[www.saferlakecounty.org](http://www.saferlakecounty.org)

